



King's Research Portal

DOI:

[10.1098/rspa.2009.0014](https://doi.org/10.1098/rspa.2009.0014)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Illingworth, C. J., & Domene, C. (2009). Many-body effects and simulations of potassium channels. *Royal Society of London. Proceedings A. Mathematical, Physical and Engineering Sciences*, 465(2106), 1701-1716. <https://doi.org/10.1098/rspa.2009.0014>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

REVIEW

Many-body effects and simulations of potassium channels

BY CHRISTOPHER J. ILLINGWORTH AND CARMEN DOMENE*

*Physical and Theoretical Chemistry Laboratory, Department of Chemistry,
University of Oxford, Oxford OX1 3QZ, UK*

The electronic polarizability of an ion or a molecule is a measure of the relative tendency of its electron cloud to be distorted from its normal shape by an electric field. On the molecular scale, in a condensed phase, any species sits in an electric field due to its neighbours, and the resulting polarization is an important contribution to the total interaction energy. Electrostatic interactions are crucial for determining the majority of chemical–physical properties of the system and electronic polarization is a fundamental component of these interactions. Thus, polarization effects should be taken into account if accurate descriptions are desired. In classical computer simulations, the forces required to drive the system are typically based on interatomic interaction potentials derived in part from electronic structure calculations or from experimental data. Owing to the difficulties in including polarization effects in classical force fields, most of them are based just on pairwise additive interaction potentials. At present, major efforts are underway to develop polarizable interaction potentials for biomolecular simulations. In this review, various ways of introducing explicit polarizability into biomolecular models and force fields are reviewed, and the progress that might be achieved in applying such methods to study potassium channels is described.

Keywords: molecular dynamics simulations; polarization effects; force fields;
ab initio; K^+ channels

1. Introduction

A force field refers to the functional form and parameter sets used to describe the potential energy of a system of particles. Force field functions and parameter sets are derived from both experimental work and high-level quantum-mechanical calculations. In particular, at the heart of the potential functions are the atomic partial charges and the Lennard-Jones radii, which surprisingly vary widely from one force field to another. The polarizability of a molecule or an ion is generally smaller in condensed phase than in gas phase because of the presence of a confining potential due to the surrounding species. The fact that the motion of every single atom influences and depends on the motion of the surrounding atoms, and therefore that coupled equations are required to describe the dynamics of the

* Author for correspondence (carmen.domene@chem.ox.ac.uk).

system, is referred as the many-body effect. Generally, most biomolecular force fields and simulations do not consider electronic polarization explicitly, using fixed permanent partial charges on the atoms or interaction sites. If polarization effects are included, it is usually in an empirical manner by a mean-field approach in which the gas-phase atomic charges are used to mimic the bulk-phase dipole moment in combination with a scaling factor reproducing many structural and dynamical properties. This represents a reasonable first-order approximation, leading to over-polarization when the ions or molecules are transferred to their condensed phases. However, in complex biological systems such as ion channels, where the ion chemical environments that should be described are very different, the mean-field approach might be inadequate. While a charge–charge interaction term coupled with Lennard-Jones forces would reproduce some of the effects present in such a system, they would not reproduce, for example, the changes in the electronic structure of carbonyls in the filter region of a potassium channel such as KcsA, which would be expected as a potassium ion proceeds through the selectivity filter region. This fact has been recognized in the development of potential functions specifically designed for ions occupying this area of the protein, which differ significantly from those for general usage in solutions (Roux & Berneche 2002). Owing to the approximations required in the creation of a classical force field, the optimal parametrization is the result of a compromise between, for example, an accurate representation of the microscopic energies and bulk solvation properties. For instance, despite its key role in biology, no universally satisfactory model potential has yet been built for liquid water (Yu & van Gunsteren 2005) even if several pairwise-additive, site–site models, such as TIP3P, TIP4P or SPC/E, have been described and extensively employed.

2. Potassium channels

Ion channels are a large and biomedically important family of integral membrane proteins (Hille 2001) that govern the electrical properties of the membranes of excitable cells such as neurons or muscle, although they are also found in the membranes of non-excitable cells and a wide range of organisms from viruses to plants. Ion channels form pores in biological membranes, and each time the channel opens, thousands of millions of ions diffuse down their electrochemical gradient across the membrane. K^+ channels are perhaps the most extensively studied family of ion channels, both experimentally and computationally. K^+ channels are selective to K^+ ions over other cations such as Na^+ . For instance, the voltage-gated KcsA channel has a K^+ flow of approximately 10^7 ions per second per channel versus only 10^4 Na^+ ions transported, even if the charge of these ions is the same, and they have very similar ionic radius (1.33 Å for K^+ and 0.95 Å for Na^+). The selectivity filter of K^+ channels is the structural element essential to the permeation and the selectivity mechanisms of these membrane protein architectures (figure 1). Ions in the extracellular and intracellular milieu cross the cellular membrane through the protein and its selectivity filter. Four identical subunits that are symmetrically disposed forming the central pore are composed of a conserved signature sequence peptide, the TVGYG motif (Thr-Val-Gly-Tyr-Gly) (Morais-Cabral *et al.* 2001; Zhou *et al.* 2001). The carbonyl oxygens of the backbone of this motif point towards the lumen, orchestrating the




Figure 1. The KirBac1.1 potassium channel (PDB 1P7B) embedded in a lipid bilayer, with individual lipids rendered as cyan chains. The transmembrane α -helices are rendered as blue ribbons and the extracellular domain is coloured according to its structure (yellow, beta sheet; white, random coil; cyan, turns). K^+ ions are rendered as purple spheres, and water oxygen and hydrogen as red and white spheres, respectively. Only waters in the selectivity filter are shown. On the right, atomic detail of the selectivity filter (TVGYG motif) in a representative conductive structure is presented. For visual clarity, atomic detail is given only of two diametrically opposed subunits.

movements of ions in and out of the channel. These key carbonyl oxygens together with the side-chain hydroxyl oxygen of the threonine residue define four equally spaced ion-binding sites. The selectivity filter is too narrow to fit an ion with its solvation shell. Thus, ions must be dehydrated to enter this region of the channel and thus K^+ ions replace the solvation shell by the carbonyl oxygens of the protein while travelling across the channel. Each of these protein sites binds the permeant cation with a tight-fitting cage of eight carbonyl oxygen atoms. This geometry resembles the solvation shell of hydrated K^+ ions. Selectivity of K^+ ions over other ions such as Na^+ was suggested to rely on the pore having a snug fit for only K^+ ions (Doyle *et al.* 1998; Gouaux & MacKinnon 2005). This suggestion would imply conformational rearrangements to discriminate two species of very close sizes. Later, it was proposed that K^+ permeation at the mouth level of the protein is facilitated owing to the fact that dehydration of K^+ ions is better compensated by reassociation with the protein carbonyls in contrast to Na^+ ions (Guidoni *et al.* 2000). More recently, it was suggested that the selectivity is determined by the ability of the channel to coordinate only some particular ion species and that the magnitude of the dipole moment of the carbonyl ligands is the predominant factor for the selectivity (Karreman & Eisenman 1962; Aqvist *et al.* 1992; Noskov *et al.* 2004). However, the dipole strength, of itself could lead to the smaller ions, Na^+ , being favoured in the pore over K^+ . Using molecular

dynamics simulations and *ab initio* calculations on small models, evidence was later presented suggesting that the magnitude of the dipole moment of the carbonyl ligands is not the predominant factor in K^+ selectivity, but that selectivity arises primarily from limitations on the way in which these dipoles can orient around an ion (Thomas *et al.* 2007; Varma & Rempe 2007; Varma *et al.* 2008). Based on quantum calculations, it was also proposed (Huetz *et al.* 2006) that a different loss of charge between K^+ and Na^+ across the channel could significantly contribute to this specific conduction and selectivity. Large polarization effects on the K^+ ions and the oxygen ligands were reflected in the values of the partial charges associated with these sets of atoms. These effects were larger for K^+ than for Na^+ , which meant that Na^+ ions are much more strongly trapped in the selectivity filter once they have managed to get in than K^+ ions. The weaker binding resulting for K^+ ions explained the higher translocation rate across the channel.

3. Approaches to polarization

Transferability of the functional form and parameters is an important feature of a force field, i.e. the same set of parameters should be capable of describing related systems. In the context of ion channel simulations, they usually do not allow for polarization effects (Roux & Berneche 2002), which is a possible limitation shared with a number of other studies of ion selectivity (Allen *et al.* 2000; Aqvist & Luzhkov 2000; Domene & Sansom 2003) and permeation (Berneche & Roux 2001; Domene *et al.* 2004; Holyoake *et al.* 2004; Furini *et al.* 2009). Certainly, electronic polarization will play a key role during the movement of the ions both in the aqueous solution and in the channel (Guidoni *et al.* 2000; Bucher *et al.* 2006) and it will merit further study.

A significant number of computational studies have been carried out on ion channels, potassium channels in particular, of which many have made use of classical force fields. These force fields, which include Chemistry at Harvard Macromolecular Mechanics (CHARMM; MacKerell *et al.* 1998), Assisted Model Building and Energy Refinement (AMBER; Cornell *et al.* 1995), GRONINGEN MOLECULAR SIMULATION (GROMOS; Oostenbrink *et al.* 2004), Optimized Potentials for Liquid Simulations–All Atom (OPLS-AA; Jorgensen & Tiradorives 1988) and many others, have a long track record of use in molecular modelling. However, the accuracy of such force fields for the simulation of ion channel systems has been called into question, for example by the work of Roux & Berneche (2002), who demonstrated the varying interaction energies between ions and *N*-methyl acetaldehyde with a range of force fields. While these authors are not entirely dismissive of the possibility for correctly calibrated molecular dynamics studies to produce meaningful results, the question is raised as to whether the use of models allowing for induced polarization might yield better data. Other authors have also raised the question of potential issues caused by the neglect of polarization (Tieleman *et al.* 2001; Domene & Sansom 2003; Jorgensen *et al.* 2007), and further studies demonstrating the importance of polarization in other ionic systems (Stuart & Berne 1996), as well as in ion channels themselves (Allen *et al.* 2004; Noskov *et al.* 2004; Bucher *et al.* 2006; Warshel *et al.* 2007), make this question all the more pertinent. As has been done in a number of cases, the use of *ab initio* methods avoids this difficulty (Guidoni *et al.* 2000;




Figure 2. Qualitative description of approaches to modelling polarization. (a) *Neglect*. Classical modelling of systems is regularly carried out with fixed atomic charges. (b) *Fluctuating charge model*. Atomic charges are modified to reproduce the effect of polarization. (c) *Drude oscillator model*. A point charge is attached to each atom by means of a spring, and moved, in order to reproduce polarization effects. (d) *Induced dipole model*. An explicit dipole term is added to each point charge. (e) *Quantum-mechanical model*. A wave function is calculated in order to model polarization on an electronic level.

Ban *et al.* 2004; Bliznyuk & Rendell 2004; Compoin *et al.* 2004; Huetz *et al.* 2006; Kariev *et al.* 2007; Kariev & Green 2008). However, *ab initio* calculations on large molecular systems are still computationally expensive and somewhat impractical. In the past, a number of alternative methods have been suggested to incorporate polarization into such molecular models, a number of which attempt to model polarization by means of alterations to a standard classical framework. These methods that are illustrated in a qualitative manner in figure 2, are often divided into four broad approaches.

(a) The fluctuating charge model

Of these approaches, the fluctuating charge method of polarization, represented in figure 2b, was developed by Rick *et al.* (1994) on the basis of previous work by Rappe & Goddard (1991), is perhaps the closest to the original concept of a classical force field. Instead of applying the fixed charges typical of such a force field, charges are allowed to vary. Where standard classical force fields model the electrostatic energy of a system as the sum of atomic charge–charge interactions, the fluctuating charge method incorporates an additional electrostatic energy term, representing the energy required to modify the charge of an individual atom. The total electrostatic energy is then minimized with respect to the individual atomic charges, subject to a set of constraints, for example keeping constant the overall charges on specific molecules, or on the system as a whole.

Specifically, as expressed by Rappe and Goddard, in a system of N atoms with charges q_1, q_2, \dots, q_N , the electrostatic energy is written as

$$E = \sum_{a=1}^N (E_{a0} + \chi_a^0 q_a) + \frac{1}{2} \sum_{a=1}^N \sum_{b=1}^N q_a q_b J_{ab}, \quad (3.1)$$

where E_{a0} is the energy of atom a at zero charge; χ_a^0 is the electronegativity of atom a ; and J_{ab} is a term equal to the Coulomb interaction between unit charges at atom locations a and b if $a \neq b$, and equal to the difference between the ionization potential and the electron affinity of the atom if $a = b$. Charges in this system are derived by calculating the N charges q_1, q_2, \dots, q_N subject to the N constraints

$$\frac{\delta E}{\delta q_1} = \frac{\delta E}{\delta q_2} = \dots = \frac{\delta E}{\delta q_N} \quad (3.2)$$

and

$$\sum_{a=1}^N q_a = Q, \quad (3.3)$$

where Q is a fixed charge for the system as a whole.

This model has found a significant number of applications to a variety of small-molecule systems, including models of water (Rick *et al.* 1994; Chen *et al.* 2000*a,b*; Rick 2001; Olano & Rick 2005), ethanol (Wang & Cann 2007) and ionic systems (Bryce *et al.* 1998; Ribeiro & Almeida 1999). More recently, the fluctuating charge model has been applied to classical force fields for modelling protein systems, being combined with the CHARMM force field (Patel *et al.* 2004), and in the work of Friesner and co-workers, being used in a stand-alone sense (Applequist *et al.* 1972; Banks *et al.* 1999), and in later work as part of a model including polarizable dipoles (Stern *et al.* 2001).

While the fluctuating charge model can be used as part of quantum mechanics/molecular mechanics (QM/MM) calculations (Field 1997) or in a stand-alone classical model (Olano & Rick 2005), the quantum mechanics/molecular mechanics (induced-charge), QMMM(IC), model of polarization developed by Reynolds and co-workers (Ferenczy *et al.* 1997; Winn *et al.* 1997, 1999; Ferenczy & Reynolds 2001; Illingworth *et al.* 2006) is explicitly designed for a QM/MM framework. Here, the charge distribution of the QM region is used to calculate induced dipoles on the MM points in the system, making use of the isotropic polarizabilities of Miller & Savchik (1979). If the field induced by the QM region is \mathbf{E}_a for some atom a , then the dipole vector $\boldsymbol{\mu}_a$ is given by the equation

$$\boldsymbol{\mu}_a = \alpha_a \mathbf{E}_a, \quad (3.4)$$

where α_a is the appropriate atomic polarizability.

The effect of these dipoles is then approximated by modifying the surrounding point charges (Ferenczy 1991). This produces a new set of point charges, which are fed back into the QM/MM model, setting up an iteration procedure that converges in a few steps to give a polarized system. In common with the fluctuating charge model, polarization is here included without the addition of dipoles or additional off-atom charges, potentially allowing for easy incorporation into modelling software. This method has successfully been applied to energy calculations in both small-molecule (Illingworth *et al.* 2006) and enzymatic (Illingworth *et al.* 2008*b*) systems, and to the problem of ligand docking (Illingworth *et al.* 2008*a*).

(b) The Drude oscillator model

The Drude oscillator model (Drude 1902), also referred to as the shell method (Dick & Overhauser 1958) or as the charge-on-spring method (Yu & van Gunsteren 2005), incorporates polarization into a classical force field by means of a set of additional massless charges attached, one to each of the atomic centres, by means of harmonic springs. Figure 2*c* gives a qualitative picture of this method. The charge at the atomic centre is correspondingly reduced by the value of the added charge, and the springs are kept short in order to recreate the effect of the original charge on the surrounding system.

Spelling this out mathematically, suppose that a massless particle with a charge of $-Q_a$ is attached to the site of an atom a , by a short spring with force constant k_a . The charge on the atom is correspondingly increased by Q_a . When the spring is in equilibrium, the force due to the electric field will be equal to the force on the spring. Thus, where the displacement of the massless particle is denoted by the vector $\Delta\mathbf{r}_a$,

$$\mathbf{E}_a Q_a = k_a \Delta\mathbf{r}_a. \quad (3.5)$$

The displacement of the massless particle creates a dipole, $\boldsymbol{\mu}_a$, in the direction of the displacement

$$\boldsymbol{\mu}_a = Q_a \Delta\mathbf{r}_a. \quad (3.6)$$

And combining equations (3.5) and (3.6), and rearranging term, gives

$$\boldsymbol{\mu}_a = \frac{(Q_a)^2}{k_a} \mathbf{E}_a. \quad (3.7)$$

Here the scalar term on the r.h.s. of the equation, equal to the square of the added charge divided by the spring constant, is analogous to the isotropic atomic polarizability of equation (3.4).

In the presence of an electric field, this addition of charged particles recreates the effect of induced dipoles at the atomic centres. As with the fluctuating charge model, this method has been applied in a significant body of work modelling water and other small-molecule systems (Sprik & Klein 1988; Saint-Martin *et al.* 2000; Hernandez-Cobos *et al.* 2005; Yu & van Gunsteren 2005; Yu *et al.* 2006; Anisimov *et al.* 2007). Considering the role of polarization in ion channels, Roux and co-workers have developed a model for water on this basis (Lamoureux *et al.* 2003, 2006), and applied it to a model of the hydration of K^+ ions (Whitfield *et al.* 2007). The Drude oscillator model was found to give results consistent with *ab initio* calculations, and in good agreement with the experimental measurements of the radial distribution function, thereby suggesting the potential of this method for application to systems of greater complexity. In addition to studies of small-molecule systems, polarization models based on the Drude oscillator model have also been applied to protein systems, being implemented in a range of common classical molecular dynamics frameworks (Lindahl *et al.* 2001; Lamoureux & Roux 2003; Yu & van Gunsteren 2004; Anisimov *et al.* 2005).

(c) Induced dipoles

A third model for polarization in classical systems incorporates explicit dipoles at each of the atomic points, illustrated by figure 2d. The dipole $\boldsymbol{\mu}_a$ at a point a with polarizability α_a is expressed as

$$\boldsymbol{\mu}_a = \alpha_a \left(\mathbf{E}_a - \sum_{\substack{b=1 \\ b \neq a}}^N T_{ab} \boldsymbol{\mu}_b \right), \quad (3.8)$$

where T_{ab} is a tensor calculated from the relative positions of the atoms a and b . Note again, the similarity between this equation and equation (3.4), and the additional term modelling the dipole–dipole interaction. The polarizability, α_a , is not necessarily isotropic, and therefore takes the form of a matrix.

Equation (3.8) is solved by rearranging into the form

$$\alpha_a^{-1} \boldsymbol{\mu}_a + \sum_{\substack{b=1 \\ b \neq a}}^N T_{ab} \boldsymbol{\mu}_b = \mathbf{E}_a, \quad (3.9)$$

$$\begin{pmatrix} \alpha_1^{-1} & T_{12} & \cdots & T_{1N} \\ T_{21} & \alpha_2^{-1} & \cdots & T_{2N} \\ \vdots & & & \vdots \\ T_{N1} & & & \alpha_N^{-1} \end{pmatrix} \begin{pmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \\ \vdots \\ \boldsymbol{\mu}_N \end{pmatrix} = \begin{pmatrix} \mathbf{E}_1 \\ \mathbf{E}_2 \\ \vdots \\ \mathbf{E}_N \end{pmatrix}, \quad (3.10)$$

where the components T_{ab} and α_a^{-1} are 3×3 matrices and the $\boldsymbol{\mu}_a$ and \mathbf{E}_a terms are 3×1 vectors. This gives a system of $3N$ equations, solved for the dipole terms $\boldsymbol{\mu}_a$, by inverting the $3N \times 3N$ matrix.

Early work employing this method was carried out by Applequist *et al.* (1972) and Warshel & Levitt (1976). Thole (1981), incorporated a damping scheme to the dipole–dipole interactions in order to avoid a situation in which dipoles placed too close together lead to an infinite polarizability. This work in turn was the basis for a series of publications including those by Ponder and co-workers (Ren & Ponder 2002, 2003; Schnieders *et al.* 2007), in which induced dipoles were used to study small-molecule systems and solvation effects. Indeed, the method of modelling polarization by explicit induced dipoles has been applied to models of a wide range of systems (Borodin & Smith 2006*a,b*; Mayer *et al.* 2006), including force fields designed for protein systems (Wang *et al.* 2006). Calculations by Baucom *et al.* (2004) have demonstrated that this brings about a meaningful improvement in the modelling of a DNA strand.

A more recent adaptation of this method replaces the framework of point charges and dipoles with Gaussian functions (Chelli & Procacci 2002; Paricaud *et al.* 2005), for example, through the use of s-orbital functions in place of point charges, and p-orbital functions in place of the point dipoles. This offers an alternative to Thole's damping method, which avoids the polarization catastrophe so long as the Gaussian functions are sufficiently diffuse. Use of such Gaussian functions has been shown to give an improved fit to molecular polarizability tensors when compared with Thole's model (Elking *et al.* 2007).

Also along these lines, the sum of interactions between fragments computed *ab initio* (SIBFA) method, developed by Gresh *et al.* (1984), is not strictly a classical method, in that it incorporated more than just polarization into the classical model, including, for example, an electrostatic term based upon *ab initio* calculations on fragments of the molecules of a system. However, it can be broadly fitted into this category, in that it makes use of point dipoles to model polarization. The SIBFA method has been used in a number of studies of ligand docking (Antony *et al.* 2005; Miller Jenkins *et al.* 2007), as well as in other applications, summarized in a recent review paper (Gresh *et al.* 2007).

The incorporation of atom-centred dipoles into a classical force field is an approximation to the higher order representation of the electrostatics of a molecule that can be achieved by means of multipole series. At the most basic level, the electrostatics of a small molecule can be represented by a single point

with fixed charge, for example

$$q = \sum_{a=1}^N q_a, \quad (3.11)$$

where q_a is the charge on atom a , in the molecule.

A more detailed description is given by adding to this a dipole term

$$\mu_i = \sum_{a=1}^N q_a a_i, \quad i = 1, 2, 3, \quad (3.12)$$

where the atom a has coordinates (a_1, a_2, a_3) .

Higher order terms can be added to the description, the next being the quadrupole term

$$\Theta_{ij} = \sum_{a=1}^N q_a \left(\frac{3}{2} a_i a_j - \frac{1}{2} a^2 \delta_{ij} \right), \quad i, j = 1, 2, 3, \quad (3.13)$$

where δ_{ij} is the Kronecker delta function. Higher order terms can also be calculated, the sum to infinity of all the terms giving a complete description of the electrostatics. While potentially useful, this approach suffers from slow convergence, a large number of terms being required to give a good description of the behaviour of a molecule.

A significant development, therefore, in the use of multipoles, was the work of Stone on the derivation of atom-centred multipoles from a charge distribution (Stone 1981). Using a set of multipole series, centred on the atoms of a molecule, led to much faster convergence, and thereby a more practical tool for calculation. This was later extended to model the response of a molecule to an external electric field (Stone 1985). More recent development of this work has included the development of distributed multipole methods for evaluating the induction energies in condensed phase of small organic molecules, a project with direct application to organic crystal structure prediction (Misquitta & Stone 2008a,b; Misquitta *et al.* 2008; Welch *et al.* 2008).

(d) QM-based polarization

A fourth method for including polarization, distinct from those above, involves the use of *ab initio* calculations on small parts of the molecules concerned, aiming to retain the accuracy of the full QM model, without carrying out an *ab initio* calculation on the whole system. Whereas the methods above broadly take the approach of adding components to a classical force field, this method takes a different ideological direction, beginning with a QM model that is simplified to be computationally tractable. An early example of this concept is the work of Field (1997) who incorporates semi-empirical molecular orbital methods into a QM/MM framework, applying the method of flexible charges to the classically modelled atoms. This approach has been taken in the MODEL potential method of Gao and co-workers (Gao 1997), in which individual molecules of a system are described at a semi-empirical level, with interactions between molecules being calculated on a QM/MM basis. This was used to develop a model for water (Gao 1998), and more recently to describe a force field suitable for calculations on peptides (Xie & Gao 2007).

In another approach, a fit is performed on a representative training set of molecules to reproduce some of their electronic properties calculated by accurate *ab initio* methods. Then the resulting atomic parameters are transferable both to new isolated molecular systems and to complexes in condensed-phase simulations (Kaminski *et al.* 2004; Chelli *et al.* 2005). A more accurate quantum-mechanical modelling of electronic polarization would be prohibitively expensive.

Simulations of biological systems often necessitate the use of large simulation cells and run times, which puts them outside the range of direct *ab initio* simulations. Potentials may then be parametrized by fitting the predicted forces and multipoles to a large body of information generated from *ab initio* calculations. The resulting potentials are predictive of *ab initio* accuracy and have a high degree of transferability (Madden *et al.* 2006). Compared with the almost parameter-free first-principles molecular dynamics in which electrons are explicitly represented, larger simulation cells and longer simulation times can be reached. A drawback although is that additional parameters are introduced in the interaction potential, and it is not always straightforward to choose them in a physically meaningful way. In the case of the polarizability, gas-phase values have often been measured experimentally, but values in liquid phase remain undetermined.

4. Discussion

Despite a considerable body of work on the problem, consensus has not yet been reached on a universally applicable model for polarization (Jorgensen 2007). Each of the classical approaches to modelling polarization is open to criticism. For example, despite recent efforts (Soteras *et al.* 2007), there is, as yet, no one agreed-upon method of defining atomic polarizabilities, essential for a dipole-based model. Drude oscillator models likewise present a question as to how to generate a correct parametrization. While if point charges alone are used to recreate polarization effects, there exists a geometrical restriction in polarization inherent to the location of the points. Although the use of additional off-atom charges provides a potential solution to this, this itself presents an issue of parametrization (Illingworth *et al.* 2006). QM-based models perhaps provide a better alternative, although they do not quite fit the ideal of incorporating polarization into a classical system.

Given the diversity of approaches summarized above, it is difficult to suggest which among them might prove the most useful for specific application to ion channels. In classical simulations where polarization is not included, there exist a number of independently derived force fields that have each produced valuable results, and, similarly, different models of polarization might each give insight into these systems. With regard to ion channel calculations, a simple modification of the ionic charge, or the addition of a charge-on-spring or dipole term to the ion itself may not be complex enough to capture the subtlety of the protein–ion and protein–water interactions, and therefore, for the ions at least, a QM-based model may be required to get the best result. If so, this would perhaps favour the use of QM-based or fluctuating charge-type approaches, these being more straightforwardly compatible with *ab initio* or QM/MM calculations. However, understanding of how best to incorporate polarization into models of these systems will most probably be gained through future experimentation.

Here we note that, while no one method of incorporating polarization into classical models has been agreed upon, there is significant agreement on the importance of polarization in ion channels, as noted above. While polarizable classical force fields have been applied to a great number of small-molecule systems, and to a variety of protein simulations, the number of applications to ion channels is relatively small. [Aqvist & Warshel \(1989\)](#) implemented a method of Langevin dipoles in order to model the solvation of sodium ions by gramicidin A. Later studies have adopted similar methods in order to model polarizability in the KcsA channel ([Luzhkov & Aqvist 2000](#); [Burykin *et al.* 2003](#)). [Bucher *et al.* \(2006\)](#) showed that there is a significant transfer of charge from the backbone atoms of the KcsA potassium channel, to the bound cations, thus the lack of polarization is a clear source of error. Given the recognized importance of polarization in these systems, ion channels perhaps provide a fertile ground for the future development and application of classical polarization techniques.

Various ways of introducing explicit polarizability into biomolecular models and force fields have been reviewed and discussed. They undoubtedly represent a significant step forward in devising new and physically realistic force fields for simulations of complex systems. However, it remains the case that all available polarizable force fields are based on classical electrostatics, none of which explicitly consider important non-classical effects such as polarization–exchange coupling or charge transfer.

Transferability of the functional form and parameters is an important feature of a force field; that is, the same set of parameters should be capable of describing related systems. In the context of ion channel simulations, they usually do not allow for polarization effects ([Roux & Berneche 2002](#)), which is a possible limitation shared with a number of other studies of ion selectivity ([Allen *et al.* 2000](#); [Aqvist & Luzhkov 2000](#)) and permeation ([Berneche & Roux 2001](#)). In the absence of a fully developed polarizable force field for ions, water, protein and lipids, the studies carried up to now have delineated some of the main features of ion channels. However, it is expected that future research directed at using polarizable force fields and QM/MM and *ab initio* simulation methods, with the advent of linear-scaling quantum-mechanical methods, will lead to further insights into these biologically important systems.

C.D. thanks The Royal Society for a University Research Fellowship. This work was supported by grants from The Leverhulme Trust and the EPSRC.

References

- Allen, T. W., Kuyucak, S. & Chung, S. H. 2000 Molecular dynamics estimates of ion diffusion in model hydrophobic and KcsA potassium channels. *Biophys. Chem.* **86**, 1–14. ([doi:10.1016/S0301-4622\(00\)00153-8](#))
- Allen, T. W., Andersen, O. S. & Roux, B. 2004 Energetics of ion conduction through the gramicidin channel. *Proc. Natl Acad. Sci. USA* **101**, 117–122. ([doi:10.1073/pnas.2635314100](#))
- Anisimov, V. M., Lamoureux, G., Vorobyov, I. V., Huang, N., Roux, B. & MacKerell, A. D. 2005 Determination of electrostatic parameters for a polarisable force field based on the classical Drude oscillator. *J. Chem. Theory Comput.* **1**, 153–168. ([doi:10.1021/ct049930p](#))
- Anisimov, V. M., Vorobyov, I. V., Roux, B. & MacKerell, A. D. 2007 Polarizable empirical force field for the primary and secondary alcohol series based on the classical Drude model. *J. Chem. Theory Comput.* **3**, 1927–1946. ([doi:10.1021/ct700100a](#))

- Antony, J., Piquemal, J. P. & Gresh, N. 2005 Complexes of thiomandelate and captopril mercaptocarboxylate inhibitors to metallo-beta-lactamase by polarisable molecular mechanics. Validation on model binding sites by quantum chemistry. *J. Comput. Chem.* **26**, 1131–1147. (doi:10.1002/jcc.20245)
- Applequist, J., Carl, J. R. & Fung, K. K. 1972 Atom dipole interaction model for molecular polarisability. Application to polyatomic molecules and determination of atom polarisabilities. *J. Am. Chem. Soc.* **94**, 2952–2960. (doi:10.1021/ja00764a010)
- Aqvist, J. & Luzhkov, V. 2000 Ion permeation mechanism of the potassium channel. *Nature* **404**, 881–884. (doi:10.1038/35009114)
- Aqvist, J. & Warshel, A. 1989 Energetics of ion permeation through membrane channels—solvation of Na^+ by gramicidin A. *Biophys. J.* **56**, 171–182. (doi:10.1016/S0006-3495(89)82662-1)
- Aqvist, J., Alvarez, O. & Eisenman, G. 1992 Ion-selective properties of a small ionophore in methanol studied by free-energy perturbation simulations. *J. Phys. Chem.* **96**, 10 019–10 025. (doi:10.1021/j100203a079)
- Ban, F. Q., Kusalik, P. & Weaver, D. F. 2004 Density functional theory investigations on the chemical basis of the selectivity filter in the K^+ channel protein. *J. Am. Chem. Soc.* **126**, 4711–4716. (doi:10.1021/ja0367290)
- Banks, J. L., Kaminski, G. A., Zhou, R. H., Mainz, D. T., Berne, B. J. & Friesner, R. A. 1999 Parametrizing a polarizable force field from *ab initio* data. I. The fluctuating point charge model. *J. Chem. Phys.* **110**, 741–754. (doi:10.1063/1.478043)
- Baucom, J., Transue, T., Fuentes-Cabrera, M., Krahn, J. M., Darden, T. A. & Sagui, C. 2004 Molecular dynamics simulations of the d(CCAACGTTGG)(2) decamer in crystal environment: comparison of atomic point-charge, extra-point, and polarisable force fields. *J. Chem. Phys.* **121**, 6998–7008. (doi:10.1063/1.1788631)
- Berneche, S. & Roux, B. 2001 Energetics of ion conduction through the K^+ channel. *Nature* **414**, 73–77. (doi:10.1038/35102067)
- Bliznyuk, A. A. & Rendell, A. P. 2004 Electronic effects in biomolecular simulations: investigation of the KcsA potassium ion channel. *J. Phys. Chem. B* **108**, 13 866–13 873. (doi:10.1021/jp0487298)
- Borodin, O. & Smith, G. D. 2006a Development of many-body polarisable force fields for Li-battery applications: 2. LiTFSI-doped oligoether, polyether, and carbonate-based electrolytes. *J. Phys. Chem. B* **110**, 6293–6299. (doi:10.1021/jp055080d)
- Borodin, O. & Smith, G. D. 2006b Development of many-body polarisable force fields for Li-battery components: 1. Ether, alkane, and carbonate-based solvents. *J. Phys. Chem. B* **110**, 6279–6292. (doi:10.1021/jp055079e)
- Bryce, R. A., Vincent, M. A., Malcolm, N. O. J., Hillier, I. H. & Burton, N. A. 1998 Cooperative effects in the structuring of fluoride water clusters: *ab initio* hybrid quantum mechanical/molecular mechanical model incorporating polarizable fluctuating charge solvent. *J. Chem. Phys.* **109**, 3077–3085. (doi:10.1063/1.476900)
- Bucher, D., Raugei, S., Guidoni, L., Dal Peraro, M., Rothlisberger, U., Carloni, P. & Klein, M. L. 2006 Polarization effects and charge transfer in the KcsA potassium channel. *Biophys. Chem.* **124**, 292–301. (doi:10.1016/j.bpc.2006.04.008)
- Burykin, A., Kato, M. & Warshel, A. 2003 Exploring the origin of the ion selectivity of the KcsA potassium channel. *Proteins* **52**, 412–426. (doi:10.1002/prot.10455)
- Chelli, R. & Procacci, P. 2002 A transferable polarisable electrostatic force field for molecular mechanics based on the chemical potential equalization principle. *J. Chem. Phys.* **117**, 9175–9189. (doi:10.1063/1.1515773)
- Chelli, R., Vincenzo Schettino, V. & Procacci, P. 2005 Comparing polarisable force fields to *ab initio* calculations reveals nonclassical effects in condensed phases. *J. Chem. Phys.* **122**, 234 107–234 114. (doi:10.1063/1.1931653)
- Chen, B., Potoff, J. J. & Siepmann, J. I. 2000a Adiabatic nuclear and electronic sampling Monte Carlo simulations in the Gibbs ensemble: application to polarisable force fields for water. *J. Phys. Chem. B* **104**, 2378–2390. (doi:10.1021/jp992459p)

- Chen, B., Xing, J. H. & Siepmann, J. I. 2000*b* Development of polarizable water force fields for phase equilibrium calculations. *J. Phys. Chem. B* **104**, 2391–2401. (doi:10.1021/jp993687m)
- Compoin, M., Ramseyer, C. & Huetz, P. 2004 *Ab initio* investigation of the atomic charges in the KcsA channel selectivity filter. *Chem. Phys. Lett.* **397**, 510–515. (doi:10.1016/j.cplett.2004.09.033)
- Cornell, W. D. *et al.* 1995 A 2nd generation force-field for the simulation of proteins, nucleic-acids, and organic molecules. *J. Am. Chem. Soc.* **117**, 5179–5197. (doi:10.1021/ja00124a002)
- Dick, B. G. & Overhauser, A. W. 1958 Theory of the dielectric constants of alkali halide crystals. *Phys. Rev.* **112**, 90–103. (doi:10.1103/PhysRev.112.90)
- Domene, C. & Sansom, M. S. P. 2003 Potassium channel, ions, and water: simulation studies based on the high resolution X-ray structure of KcsA. *Biophys. J.* **85**, 2787–2800. (doi:10.1016/S0006-3495(03)74702-X)
- Domene, C., Grottesi, A. & Sansom, M. S. P. 2004 Filter flexibility and distortion in a bacterial inward rectifier K⁺ channel: simulation studies of KirBac1.1. *Biophys. J.* **87**, 256–267. (doi:10.1529/biophysj.104.039917)
- Doyle, D. A., Cabral, J. M., Pfuetzner, R. A., Kuo, A. L., Gulbis, J. M., Cohen, S. L., Chait, B. T. & MacKinnon, R. 1998 The structure of the potassium channel: molecular basis of K⁺ conduction and selectivity. *Science* **280**, 69–77. (doi:10.1126/science.280.5360.69)
- Drude, P. 1902 *The theory of optics*. New York, NY: Longmans, Green and Co.
- Elking, D., Darden, T. & Woods, R. J. 2007 Gaussian induced dipole polarization model. *J. Comput. Chem.* **28**, 1261–1274. (doi:10.1002/jcc.20574)
- Ferenczy, G. G. 1991 Charges derived from distributed multipole series. *J. Comput. Chem.* **12**, 913–917. (doi:10.1002/jcc.540120802)
- Ferenczy, G. G. & Reynolds, C. A. 2001 Modeling polarization through induced atomic charges. *J. Phys. Chem. A* **105**, 11 470–11 479. (doi:10.1021/jp0117967)
- Ferenczy, G. G., Winn, P. J. & Reynolds, C. A. 1997 Toward improved force fields. 2. Effective distributed multipoles. *J. Phys. Chem. A* **101**, 5446–5455. (doi:10.1021/jp9712011)
- Field, M. J. 1997 Hybrid quantum mechanical molecular mechanical fluctuating charge models for condensed phase simulations. *Mol. Phys.* **91**, 835–845. (doi:10.1080/002689797170941)
- Furini, S., Beckstein, O. & Domene, C. 2009 Permeation of water through the KcsA K⁺ channel. *Proteins* **74**, 437–448. (doi:10.1002/prot.22163)
- Gao, J. L. 1997 Toward a molecular orbital derived empirical potential for liquid simulations. *J. Phys. Chem. B* **101**, 657–663. (doi:10.1021/jp962833a)
- Gao, J. L. 1998 A molecular-orbital derived polarization potential for liquid water. *J. Chem. Phys.* **109**, 2346–2354. (doi:10.1063/1.476802)
- Gouaux, E. & MacKinnon, R. 2005 Principles of selective ion transport in channels and pumps. *Science* **310**, 1461–1465. (doi:10.1126/science.1113666)
- Gresh, N., Claverie, P. & Pullman, A. 1984 Theoretical studies of molecular conformation derivation of an additive procedure for the computation of intramolecular interaction energies. Comparison with *ab initio* SCF computations. *Theor. Chim. Acta* **66**, 1–20. (doi:10.1007/BF00577135)
- Gresh, N., Cisneros, G. A., Darden, T. A. & Piquemal, J. P. 2007 Anisotropic, polarizable molecular mechanics studies of inter- and intramolecular interactions and ligand–macromolecule complexes. A bottom-up strategy. *J. Chem. Theory Comput.* **3**, 1960–1986. (doi:10.1021/ct700134r)
- Guidoni, L., Torre, V. & Carloni, P. 2000 Water and potassium dynamics inside the KcsA K⁺ channel. *FEBS Lett.* **477**, 37–42. (doi:10.1016/S0014-5793(00)01712-9)
- Hernandez-Cobos, J., Saint-Martin, H., Mackie, A. D., Vega, L. F. & Ortega-Blake, I. 2005 Water liquid–vapor equilibria predicted by refined *ab initio* derived potentials. *J. Chem. Phys.* **123**, 44 506–44 508. (doi:10.1063/1.1950627)
- Hille, B. 2001 *Ionic channels of excitable membranes*. Sunderland, MA: Sinauer Associates, Inc.
- Holyoake, J., Domene, C., Bright, J. N. & Sansom, M. S. P. 2004 KcsA closed and open: modeling and simulations studies. *Eur. Biophys. J.* **33**, 238–246. (doi:10.1007/s00249-003-0355-2)
- Huetz, P., Boiteux, C., Compoin, M., Ramseyer, C. & Girardet, C. 2006 Incidence of partial charges on ion selectivity in potassium channels. *J. Chem. Phys.* **124**, 44 703. (doi:10.1063/1.2159483)

- Illingworth, C. J. R., Gooding, S. R., Winn, P. J., Jones, G. A., Ferenczy, G. G. & Reynolds, C. A. 2006 Classical polarization in hybrid QM/MM methods. *J. Phys. Chem. A* **110**, 6487–6497. (doi:10.1021/jp046944i)
- Illingworth, C. J. R., Morris, G. M., Parkes, K. E. B., Snell, C. R. & Reynolds, C. A. 2008*a* Assessing the role of polarisation in docking. *J. Phys. Chem. A* **112**, 12 157–12 163. (doi:10.1021/jp710169m)
- Illingworth, C. J. R., Parkes, K. E., Snell, C. R., Marti, S., Moliner, V. & Reynolds, C. A. 2008*b* The effect of MM polarization on the QM/MM transition state stabilization: application to chorismate mutase. *Mol. Phys.* **106**, 1511–1515. (doi:10.1080/00268970802077850)
- Jorgensen, W. L. 2007 Special issue on polarisation. *J. Chem. Theory Comput.* **3**, 1877. (doi:10.1021/ct700252g)
- Jorgensen, W. L. & Tiradorives, J. 1988 The OPLS potential functions for proteins, energy minimisations for crystals of cyclic-peptides and crambin. *J. Am. Chem. Soc.* **110**, 1657–1666. (doi:10.1021/ja00214a001)
- Jorgensen, W. L., Jensen, K. P. & Alexandrova, A. N. 2007 Polarization effects for hydrogen-bonded complexes of substituted phenols with water and chloride ion. *J. Chem. Theory Comput.* **3**, 1987–1992. (doi:10.1021/ct7001754)
- Kaminski, G. A., Stern, H. A., Berne, B. J. & Friesner, R. A. 2004 Development of an accurate and robust polarizable molecular mechanics force field from *ab initio* quantum chemistry. *J. Phys. Chem. A* **108**, 621–627. (doi:10.1021/jp0301103)
- Kariev, A. M. & Green, M. E. 2008 Quantum mechanical calculations on selectivity in the KcsA channel: the role of the aqueous cavity. *J. Phys. Chem. B* **112**, 1293–1298. (doi:10.1021/jp076854o)
- Kariev, A. M., Znamenskiy, V. S. & Green, M. E. 2007 Quantum mechanical calculations of charge effects on gating the KcsA channel. *Biochim. Biophys. Acta* **1768**, 1218–1229. (doi:10.1016/j.bbamem.2007.01.021)
- Karreman, G. & Eisenman, G. 1962 Electrical potentials and ionic fluxes in ion exchangers. I. n-Type non-ideal systems with zero current. *Bull. Math. Biophys.* **24**, 413–427. (doi:10.1007/BF02477998)
- Lamoureux, G. & Roux, B. 2003 Modeling induced polarization with classical drude oscillators: theory and molecular dynamics simulation algorithm. *J. Chem. Phys.* **119**, 3025–3039. (doi:10.1063/1.1589749)
- Lamoureux, G., MacKerell, A. D. & Roux, B. 2003 A simple polarisable model of water based on classical drude oscillators. *J. Chem. Phys.* **119**, 5185–5197. (doi:10.1063/1.1598191)
- Lamoureux, G., Harder, E., Vorobyov, I. V., Roux, B. & MacKerell, A. D. 2006 A polarisable model of water for molecular dynamics simulations of biomolecules. *Chem. Phys. Lett.* **418**, 245–249. (doi:10.1016/j.cplett.2005.10.135)
- Lindahl, E., Hess, B. & van der Spoel, D. 2001 GROMACS 3.0: a package for molecular simulation and trajectory analysis. *J. Mol. Model.* **7**, 306–317. (doi:10.1007/s008940100045)
- Luzhkov, V. B. & Åqvist, J. 2000 A computational study of ion binding and protonation states in the KcsA potassium channel. *Biochim. Biophys. Acta* **1481**, 360–370. (doi:10.1016/S0167-4838(00)00183-7)
- MacKerell, A. D. *et al.* 1998 All-atom empirical potential for molecular modeling and dynamics studies of proteins. *J. Phys. Chem. B* **102**, 3586–3616. (doi:10.1021/jp973084f)
- Madden, P. A., Heaton, R., Aguado, A. & Jahn, S. 2006 From first-principles to material properties. *J. Mol. Struct. Theochem.* **771**, 9–18. (doi:10.1016/j.theochem.2006.03.015)
- Mayer, A., Lambin, P. & Langlet, R. 2006 Charge-dipole model to compute the polarization of fullerenes. *Appl. Phys. Lett.* **89**, 167–168. (doi:10.1063/1.2337524)
- Miller, K. J. & Savchik, J. A. 1979 New empirical-method to calculate average molecular polarisabilities. *J. Am. Chem. Soc.* **101**, 7206–7213. (doi:10.1021/ja00518a014)
- Miller Jenkins, L. M., Hara, T., Durell, S. R., Hayashi, R., Inman, J. K., Piquemal, J.-P., Gresh, N. & Appella, E. 2007 Specificity of acyl transfer from 2-mercaptobenzamide thioesters to the HIV-1 nucleocapsid protein. *J. Am. Chem. Soc.* **129**, 11 067–11 078. (doi:10.1021/ja071254o)

- Misquitta, A. J. & Stone, A. J. 2008*a* Accurate induction energies for small organic molecules: 1. Theory. *J. Chem. Theory Comput.* **4**, 7–18. (doi:10.1021/ct700104t)
- Misquitta, A. J. & Stone, A. J. 2008*b* Dispersion energies for small organic molecules: first row atoms. *Mol. Phys.* **106**, 1631–1643. (doi:10.1080/00268970802258617)
- Misquitta, A. J., Stone, A. J. & Price, S. L. 2008 Accurate induction energies for small organic molecules. 2. Development and testing of distributed polarizability models against SAPT(DFT) energies. *J. Chem. Theory Comput.* **4**, 19–32. (doi:10.1021/ct700105f)
- Morais-Cabral, J. H., Zhou, Y. F. & MacKinnon, R. 2001 Energetic optimization of ion conduction rate by the K⁺ selectivity filter. *Nature* **414**, 37–42. (doi:10.1038/35102000)
- Noskov, S. Y., Berneche, S. & Roux, B. 2004 Control of ion selectivity in potassium channels by electrostatic and dynamic properties of carbonyl ligands. *Nature* **431**, 830–834. (doi:10.1038/nature02943)
- Olano, L. R. & Rick, S. W. 2005 Fluctuating charge normal modes: an algorithm for implementing molecular dynamics simulations with polarisable potentials. *J. Comput. Chem.* **26**, 699–707. (doi:10.1002/jcc.20212)
- Oostenbrink, C., Villa, A., Mark, A. E. & Van Gunsteren, W. F. 2004 A biomolecular force field based on the free enthalpy of hydration and solvation: the GROMOS force-field parameter sets 53A5 and 53A6. *J. Comput. Chem.* **25**, 1656–1676. (doi:10.1002/jcc.20090)
- Paricaud, P., Predota, M., Chialvo, A. A. & Cummings, P. T. 2005 From dimer to condensed phases at extreme conditions: accurate predictions of the properties of water by a Gaussian charge polarisable model. *J. Chem. Phys.* **122**, 244 511–244 514. (doi:10.1063/1.1940033)
- Patel, S., Mackerell, A. D. & Brooks, C. L. 2004 CHARMM fluctuating charge force field for proteins: protein/solvent properties from molecular dynamics simulations using a nonadditive electrostatic model. *J. Comput. Chem.* **25**, 1504–1514. (doi:10.1002/jcc.20077)
- Rappe, A. K. & Goddard, W. A. 1991 Charge equilibration for molecular dynamics simulations. *J. Phys. Chem.* **95**, 3358–3363. (doi:10.1021/j100161a070)
- Ren, P. Y. & Ponder, J. W. 2002 Consistent treatment of inter- and intramolecular polarisation in molecular mechanics calculations. *J. Comput. Chem.* **23**, 1497–1506. (doi:10.1002/jcc.10127)
- Ren, P. Y. & Ponder, J. W. 2003 Polarizable atomic multipole water model for molecular mechanics simulation. *J. Phys. Chem. B* **107**, 5933–5947. (doi:10.1021/jp027815+)
- Ribeiro, M. C. C. & Almeida, L. C. J. 1999 Fluctuating charge model for polyatomic ionic systems: a test case with diatomic anions. *J. Chem. Phys.* **110**, 11 445–11 448. (doi:10.1063/1.479085)
- Rick, S. W. 2001 Simulations of ice and liquid water over a range of temperatures using the fluctuating charge model. *J. Chem. Phys.* **114**, 2276–2283. (doi:10.1063/1.1336805)
- Rick, S. W., Stuart, S. J. & Berne, B. J. 1994 Dynamical fluctuating charge force-fields: application to liquid water. *J. Chem. Phys.* **101**, 6141–6156. (doi:10.1063/1.468398)
- Roux, B. & Berneche, S. 2002 On the potential functions used in molecular dynamics simulations of ion channels. *Biophys. J.* **82**, 1681–1684. (doi:10.1016/S0006-3495(02)75520-3)
- Saint-Martin, H., Hernandez-Cobos, J., Bernal-Uruchurtu, M. I., Ortega-Blake, I. & Berendsen, H. J. C. 2000 A mobile charge densities in harmonic oscillators (MCDHO) molecular model for numerical simulations: the water–water interaction. *J. Chem. Phys.* **113**, 10 899–10 912. (doi:10.1063/1.1324711)
- Schnieders, M. J., Baker, N. A., Ren, P. Y. & Ponder, J. W. 2007 Polarizable atomic multipole solutes in a Poisson–Boltzmann continuum. *J. Chem. Phys.* **126**, 124 114. (doi:10.1063/1.2714528)
- Soteras, I., Curutchet, C., Bidon-Chanal, A., Dehez, F., Angyan, J. G., Orozco, M., Chipot, C. & Luque, F. J. 2007 Derivation of distributed models of atomic polarisability for molecular simulations. *J. Chem. Theory Comput.* **3**, 1901–1913. (doi:10.1021/ct7001122)
- Språk, M. & Klein, M. L. 1988 A polarisable model for water using distributed charge sites. *J. Chem. Phys.* **89**, 7556–7560. (doi:10.1063/1.455722)
- Stern, H. A., Rittner, F., Berne, B. J. & Friesner, R. A. 2001 Combined fluctuating charge and polarizable dipole models: application to a five-site water potential function. *J. Chem. Phys.* **115**, 2237–2251. (doi:10.1063/1.1376165)

- Stone, A. J. 1981 Distributed multipole analysis, or how to describe a molecular charge-distribution. *Chem. Phys. Lett.* **83**, 233–239. (doi:10.1016/0009-2614(81)85452-8)
- Stone, A. J. 1985 Distributed polarisabilities. *Mol. Phys.* **56**, 1065–1082. (doi:10.1080/00268978500102901)
- Stuart, S. J. & Berne, B. J. 1996 Effects of polarizability on the hydration of the chloride ion. *J. Phys. Chem.* **100**, 11 934–11 943. (doi:10.1021/jp961076d)
- Thole, B. T. 1981 Molecular polarisabilities calculated with a modified dipole interaction. *Chem. Phys.* **59**, 341–350. (doi:10.1016/0301-0104(81)85176-2)
- Thomas, M., Jayatilaka, D. & Corry, B. 2007 The predominant role of coordination number in potassium channel selectivity. *Biophys. J.* **93**, 2635–2643. (doi:10.1529/biophysj.107.108167)
- Tieleman, D. P., Biggin, P. C., Smith, G. R. & Sansom, M. S. P. 2001 Simulation approaches to ion channel structure–function relationships. *Q. Rev. Biophys.* **34**, 473–561. (doi:10.1017/s0033583501003729)
- Varma, S. & Rempe, S. B. 2007 Tuning ion coordination architectures to enable selective partitioning. *Biophys. J.* **93**, 1093–1099. (doi:10.1529/biophysj.107.107482)
- Varma, S., Sabo, D. & Rempe, S. B. 2008 K^+/Na^+ selectivity in K^+ channels and valinomycin: over-coordination versus cavity-size constraints. *J. Mol. Biol.* **376**, 13–22. (doi:10.1016/j.jmb.2007.11.059)
- Wang, S. H. & Cann, N. M. 2007 Polarizable and flexible model for ethanol. *J. Chem. Phys.* **126**, 214 502. (doi:10.1063/1.2730837)
- Wang, Z. X., Zhang, W., Wu, C., Lei, H. X., Cieplak, P. & Duan, Y. 2006 Strike a balance: optimization of backbone torsion parameters of amber polarizable force field for simulations of proteins and peptides. *J. Comput. Chem.* **27**, 781–790. (doi:10.1002/jcc.20386)
- Warshel, A. & Levitt, M. 1976 Theoretical studies of enzymatic reactions—dielectric, electrostatics and steric stabilization of carbonium-ion in reaction of lysozyme. *J. Mol. Biol.* **103**, 227–249. (doi:10.1016/0022-2836(76)90311-9)
- Warshel, A., Kato, M. & Pisiakov, A. V. 2007 Polarizable force fields: history, test cases, and prospects. *J. Chem. Theory Comput.* **3**, 2034–2045. (doi:10.1021/ct700127w)
- Welch, G. W. A., Karamertzanis, P. G., Misquitta, A. J., Stone, A. J. & Price, S. L. 2008 Is the induction energy important for modeling organic crystals? *J. Chem. Theory Comput.* **4**, 522–532. (doi:10.1021/ct700270d)
- Whitfield, T. W., Varma, S., Harder, E., Lamoureux, G., Rempe, S. B. & Roux, B. 2007 Theoretical study of aqueous solvation of K^+ comparing *ab initio*, polarizable, and fixed-charge models. *J. Chem. Theory Comput.* **3**, 2068–2082. (doi:10.1021/ct700172b)
- Winn, P. J., Ferenczy, G. G. & Reynolds, C. A. 1997 Toward improved force fields. 1. Multipole-derived atomic charges. *J. Phys. Chem. A* **101**, 5437–5445. (doi:10.1021/jp971202t)
- Winn, P. J., Ferenczy, G. G. & Reynolds, C. A. 1999 Toward improved force fields: polarisation through modified atomic charges. *J. Comp. Chem.* **20**, 704–712. (doi:10.1002/(SICI)1096-987X(199905)20:7<704::AID-JCC5>3.0.CO;2-T)
- Xie, W. S. & Gao, J. L. 2007 Design of a next generation force field: the X-POL potential. *J. Chem. Theory Comput.* **3**, 1890–1900. (doi:10.1021/ct700167b)
- Yu, H. B. & van Gunsteren, W. F. 2004 Charge-on-spring polarizable water models revisited: from water clusters to liquid water to ice. *J. Chem. Phys.* **121**, 9549–9564. (doi:10.1063/1.1805516)
- Yu, H. B. & van Gunsteren, W. F. 2005 Accounting for polarization in molecular simulation. *Comput. Phys. Commun.* **172**, 69–85. (doi:10.1016/j.cpc.2005.01.022)
- Yu, H. B., Geerke, D. P., Liu, H. Y. & van Gunsteren, W. E. 2006 Molecular dynamics simulations of liquid methanol and methanol–water mixtures with polarizable models. *J. Comput. Chem.* **27**, 1494–1504. (doi:10.1002/jcc.20429)
- Zhou, Y. F., Morais-Cabral, J. H., Kaufman, A. & MacKinnon, R. 2001 Chemistry of ion coordination and hydration revealed by a K^+ channel-Fab complex at 2.0 angstrom resolution. *Nature* **414**, 43–48. (doi:10.1038/35102009)